

Gravitational Wave Background from Phantom Superinflation

Yun-Song Piao^{a,b}

^aCollege of Physical Sciences, Graduate School of Chinese Academy of Sciences, YuQuan Road 19A, Beijing 100049, China and

^bInterdisciplinary Center of Theoretical Studies, Chinese Academy of Sciences, P.O. Box 2735, Beijing 100080, China

Recently, the early superinflation driven by phantom field has been proposed and studied. The detection of primordial gravitational wave is an important means to know the state of very early universe. In this brief report we discuss in detail the gravitational wave background excited during the phantom superinflation.

PACS numbers: 98.80.Cq

Now a number of detectors for gravitational waves are (or expected to start) operating, one of whose aim is searching a stochastic background of gravitational wave. This background is expected to have different components with different origins. The primordial gravitational wave among them will be especially interesting, since it would carry information about the state of the very early universe. The basal mechanism of generation of primordial gravitational wave in cosmology has been discussed in Ref. [1, 2], see [3, 4, 5, 6, 7, 8] for the stochastic gravitational wave generated during the inflation, [9, 10, 11] for that of Pre Big Bang scenario, [12] for that of cyclic model.

Recently, the superinflation driven by the phantom field, in which the parameter of state equation $w < -1$ and the null energy condition is violated¹, has been proposed and studied in Ref. [14, 15, 16, 17, 18, 19]. The initial perturbation during the phantom superinflation will leave the horizon, and can reenter the horizon during the radiation/matter domination, which may be regarded as our late time observable universe. The phantom superinflation model leads to a blue or strong blue spectrum of primordial tensor perturbations, which is different from a nearly scale invariant (slightly red) spectrum predicted by the inflation model. The reason is that the phantom superinflation involves an early phase of super acceleration in which the Hubble parameter increases with the time, while the inflation does not. Thus it may be essential to revisit in detail the gravitational wave background from the phantom superinflation.

We set $m_p^2 = 1$ and work with the parameter $\epsilon \equiv -\dot{h}/h^2$ being constant for simplicity, where $h \equiv \dot{a}/a$. We firstly briefly show the model-independent characters of phantom superinflation. The evolution of scale factor during the phantom phase can be simply taken as $a(t) \sim (-t)^n$ [20], where t is from $-\infty$ to 0_- , and n is a negative constant. Thus we have $\epsilon = 1/n < 0$. In the conformal time, we obtain $-\eta \sim (-t)^{-n+1}$, and thus

$$a(\eta) \sim (-\eta)^{\frac{n}{1-n}} \equiv (-\eta)^{\frac{1}{\epsilon-1}}; \quad (1)$$

$$h \equiv \frac{a'}{a^2} = \frac{1}{(\epsilon-1)a\eta}, \quad (2)$$

where the prime denotes the derivative with respect to η . The perturbations leaving the horizon during the phantom superinflation can reenter the horizon during radiation/matter domination, which may be responsible for the structure formation of our observable universe. Further it may be convenient to define

$$\mathcal{N} \equiv \ln\left(\frac{a_e h_e}{a_i h_i}\right), \quad (3)$$

which measures the efolding number that the perturbation with the present horizon scale leaves the horizon before the end of the phantom phase, where the subscript e and i denote the end time of the phantom superinflation and the time that the perturbation with the present horizon scale leaves the horizon, respectively, and thus $a_i h_i = a_0 h_0$, where the subscript 0 denotes the present time, see Ref. [21]. From (1) and (2), we obtain

$$a \sim \left(\frac{1}{(1-\epsilon)ah}\right)^{\frac{1}{\epsilon-1}}. \quad (4)$$

Thus we have

$$\frac{a_e}{a_i} = \left(\frac{a_i h_i}{a_e h_e}\right)^{\frac{1}{\epsilon-1}} = e^{\frac{\mathcal{N}}{1-\epsilon}}. \quad (5)$$

We can see that during the phantom superinflation, the change of scale factor is dependent on ϵ . For the negative enough ϵ , the change $\Delta a/a = (a_e - a_i)/a_i \simeq \mathcal{N}/(1-\epsilon)$ of a can be very small. Taking the logarithm in both sides of (4), we obtain²

$$\ln\left(\frac{1}{ah}\right) = (\epsilon-1) \ln a. \quad (6)$$

We plot Fig.1 to further illustrate the characters of the phantom superinflation. We assume, throughout this

¹ Note that it is also possible to get phantom energy in scalar-tensor theories of gravity without non canonical kinetic terms [13].

² The equation can be shown and actually also applied for the expansion with arbitrary constant ϵ .




FIG. 1: The sketch of evolution of $\ln(1/ah)$ with respect to the scale factor $\ln a$ during the phantom superinflation for different ϵ , in which in the phantom superinflation region the solid line denotes the usual inflation with $\epsilon \simeq 0$, and the dot, two dots and three dots -dashed lines denote the superinflation with $\epsilon = -1, -5, -\infty$, respectively. The perturbation modes leave the Hubble horizon during the phantom superinflation and then reenter the horizon during the radiation/matter domination at late time.

brief report, that after the phantom phase ends, the reheating will rapidly occur and then bring the universe back to the usual FRW evolution.

We will discuss the primordial tensor perturbations from the phantom superinflation in the following. In the momentum space, the equation of motion of gauge invariant variable u_k , which is related to the tensor perturbation by $u_k \equiv ah_k$, is

$$u_k'' + (k^2 - \frac{a''}{a})u_k = 0, \quad (7)$$

where $a(\eta)$ is given by (1). The general solutions of this equation are the Hankel functions. In the regime $k\eta \rightarrow \infty$, all interesting modes are very deep in the horizon of phantom phase, thus (7) can be reduced to the equation of a simple harmonic oscillator $u_k \sim \frac{e^{-ik\eta}}{(2k)^{1/2}}$, which in some sense suggests that the initial condition can be taken as usual Minkowski vacuum. In the super-horizon scale, in which the modes become unstable and grow, the expansion of Hankel functions to the leading term of k gives

$$u_k \simeq \frac{1}{\sqrt{2k}}(-k\eta)^{\frac{1}{2}-\nu}, \quad (8)$$

where the phase factor and the constant with order one have been neglected, and

$$\nu^2 \equiv \frac{1}{4}(\frac{\epsilon-3}{\epsilon-1})^2. \quad (9)$$

In the phantom phase in which $\epsilon < 0$, we have $1/2 < \nu < 3/2$.

The amplitude of gravitational wave after the end of phantom superinflation is given by, accounting for both

polarizations,

$$\mathcal{P}_t^{1/2}(k, \eta_e) = \frac{k^{3/2}}{\pi} \left| \frac{u_k(-k\eta_e)}{a_e} \right| \simeq \frac{(-\eta_e)^{1/2-\nu}}{a_e} k^{3/2-\nu}. \quad (10)$$

We obtain, from (2) and (3),

$$\begin{aligned} \mathcal{P}_t^{1/2}(k, \eta_e) &= \frac{1}{\pi a_e} \left(\frac{1}{(1-\epsilon)a_e h_e} \right)^{1/2-\nu} k^{3/2-\nu} \\ &= \frac{h_e}{\pi} \frac{1}{(1-\epsilon)^{1/2-\nu}} \left(\frac{k_0}{k_e} \right)^{3/2-\nu} \left(\frac{k}{k_0} \right)^{3/2-\nu} \\ &= \frac{h_e}{\pi} \frac{e^{-\mathcal{N}(3/2-\nu)}}{(1-\epsilon)^{1/2-\nu}} \left(\frac{k}{k_0} \right)^{3/2-\nu}. \end{aligned} \quad (11)$$

We can see that for fixed $\mathcal{N}$ and h_e , when $\nu \simeq 3/2$, which corresponds to $\epsilon \simeq 0$, $\mathcal{P}_t^{1/2}(k, \eta_e) \simeq h_e/\pi$ is scale invariant, which is the usual result of inflation models, while when $\nu < 3/2$, which corresponds to $\epsilon < 0$, the spectrum of $\mathcal{P}_t^{1/2}(k, \eta_e)$ is blue tilted and has the exponentially suppressed amplitude $\sim h_e e^{-\mathcal{N}(3/2-\nu)}/\pi$ at the largest scale $k \simeq k_0$. The limit of $\nu \rightarrow 1/2$ is the solution of Ref. [14], in which the scale factor is nearly unchanged and the Hubble parameter h experiences an instantaneous “jump”, which results in that the spectrum is strong blue tilt

$$\mathcal{P}_t^{1/2}(k, \eta_e) \simeq h_e e^{-\mathcal{N}} (k/k_0)/\pi. \quad (12)$$

To convert from the primordial spectrum to the present spectrum, we need to know the transfer function $T(k)$,

$$\mathcal{P}_t^{1/2}(k, \eta_0) = T(k) \mathcal{P}_t^{1/2}(k, \eta_e). \quad (13)$$

In general, $\mathcal{P}(k, \tau)$ is roughly time independent outside the horizon and decay as a^{-1} as long as the corresponding mode reenters the horizon. Therefore, based on the fact that $h \sim a^{-2}$ during the radiation domination and $h \sim a^{-3/2}$ during the matter domination, the numerical fitting gave [22, 23]

$$T(k) \simeq \left(\frac{k_0}{k} \right)^2 \left(1 + \frac{4}{3} \frac{k}{k_{eq}} + \frac{5}{2} \left(\frac{k}{k_{eq}} \right)^2 \right)^{1/2}, \quad (14)$$

where $k_{eq} = a_{eq} h_{eq}$ denotes the modes entering the horizon at the time of matter-radiation equality. Eq. (14) actually only applies for $k > k_0$ and below this $T(k) = 1$. We have, from (5),

$$\frac{k_{eq}}{k_0} = \frac{a_{eq} h_{eq}}{a_0 h_0} = \left(\frac{a_0}{a_{eq}} \right)^{\epsilon-1}. \quad (15)$$

For the matter domination, $\epsilon = 3/2$. Thus $k_{eq}/k_0 = \sqrt{1+z_{eq}}$.

To characterize the spectrum of the stochastic gravitational wave signal, an useful quantity $\Omega_{gw}(k)$ is introduced, see [24, 25, 26], which is the gravitational wave energy per unit logarithmic wave number in the units of the critical density

$$\Omega_{gw}(k) = \frac{k}{\rho_{cr}} \frac{d\rho_{gw}}{dk}, \quad (16)$$




FIG. 2: The figure of $\Omega(k)$ from the phantom superinflation with different ϵ , where the horizon axis is k/k_0 taken from 1 to 10^{25} . We take $k_0 = 10^{-18}$ and $h_e^2 = 10^{-12}$. Thus $k_{max} \sim 10^8$ and $k_{max}/k_0 \sim 10^{26}$.

where $\rho_{cr} = 3h_0^2/8\pi$ is the critical energy density of observable universe and ρ_{gw} is the energy density of gravitational wave. The value of h_0 is generally written as $h_0 = \dot{h}_0 \times 100 \text{ km/sec/Mpc}$, where $\dot{h}_0$ parameterizes the existing experimental uncertainty. However, for simplicity we will neglect $\dot{h}_0$ in the following. In general [23],

$$\begin{aligned}\Omega_{gw}(k) &\simeq \frac{1}{6} \left(\frac{k}{k_0}\right)^2 \mathcal{P}_t(k, \eta_0) \\ &\simeq \frac{1}{6} \left(\frac{k_0}{k}\right)^2 \left(1 + \frac{4}{3} \frac{k}{k_{eq}} + \frac{5}{2} \left(\frac{k}{k_{eq}}\right)^2\right) \mathcal{P}_t(k, \eta_e)\end{aligned}\quad (17)$$

where (14) has been used. Fig.2 shows the results of $\Omega_{gw}(k)$ for the phantom superinflation with different ϵ . The spectrum is cut off at the wave number $k_{max} = k_e$. For the instantaneous reheating in which the phantom superinflation is immediately followed by a phase of radiation domination, $k_{max} = a_e h_e \simeq a_0 T_0 h_e^{1/2}$, where T_0 is the temperature of present CMB. The overall amplitude $\sim h_e^2$ of $\Omega_{gw}(k)$ should be given by the CMB observation at the large angular scale. The present bound is $\Omega_{gw}(k) < 10^{-11} k_0/k$ at $10^{-18} \text{ Hz} < k < 10^{-16} \text{ Hz}$ [27]. We can see that for $\epsilon \simeq 0$, $\Omega_{gw}(k)$ decays with k up to $k \simeq k_{eq}$, and after $k \gg k_{eq}$ it enters a nearly long plateau

$$\Omega_{gw}(k \gg k_{eq}, \epsilon \simeq 0) \simeq \frac{5h_e^2}{12\pi^2} \left(\frac{k_0}{k_{eq}}\right)^2, \quad (18)$$

which is the character of inflation model [23], while for $\epsilon < 0$ the case is distinctly different, in which at $k \gg k_{eq}$, $\Omega_{gw}(k)$ will increase with k up to $k \simeq k_{max}$. The slope of increasing is very dependent of ϵ , and the extreme one is $\sim k^2$ corresponding to $\epsilon \rightarrow -\infty$ [14]. However, since at low frequency region, the amplitude of $\Omega_{gw}(k)$ has a strong suppression from the efolding number, see (11), which is relevant with ϵ , at the case that the reheating temperature $T_r \simeq h_e^{1/2}$ is same we can obtain at $k \gg k_{eq}$

$$\begin{aligned}\Omega_{gw}(k_{max}) &\simeq (1 - \epsilon)^{2\nu-1} \frac{5h_e^2}{12\pi^2} \left(\frac{k_0}{k_e}\right)^2 \left(\frac{k_e}{k_{eq}}\right)^2 \left(\frac{k_0}{k_e}\right)^{3-2\nu} \left(\frac{k_e}{k_0}\right)^{3-2\nu} \\ &= (1 - \epsilon)^{2\nu-1} \frac{5h_e^2}{12\pi^2} \left(\frac{k_0}{k_{eq}}\right)^2 \\ &\simeq \Omega_{gw}(k \gg k_{eq}, \epsilon \simeq 0).\end{aligned}\quad (19)$$

Thus though for $\epsilon < 0$ there is an interesting increasing of $\Omega_{gw}(k)$ amplitude at high k , when the scale h_e (or reheating scale) is taken as same, the largest value obtained at $k \simeq k_{max} = k_e$ is always around the plateau $\Omega_{gw}(k \gg k_{eq}, \epsilon \simeq 0)$ of the inflation. This result is not dependent on the value of the reheating scale.

In summary, we study in detail the gravitational wave background excited during the phantom superinflation. For $\epsilon \simeq 0_-$, the amplitude of spectrum is very close to that of inflation, and can be expected to be detected in the future. But for $\epsilon \ll -1$, the amplitude is very low at the large scale (low k) and hardly seen, however, in high frequency region the amplitude increases $\sim k^{3-2\nu} = k^{2\epsilon/(\epsilon-1)}$ up to the plateau of gravitational wave background of the inflation, thus there may be some significant and different signals around k_{max} . These results may be interesting for the present and planned experiments detecting gravitational wave.

Acknowledgments This work was partly completed in Interdisciplinary Center of Theoretical Studies, Chinese Academy of Sciences, and also in part by NNSFC under Grant No: 10405029, 90403032, and National Basic Research Program of China under Grant No: 2003CB716300.

-
- [1] L.P. Grishchuk, Sov. Phys. JETP **40**, 409 (1975).
 - [2] B. Allen, Phys. Rev. **D37**, 2078 (1988).
 - [3] A. Starobinski, JETP Lett. **30**, 682 (1979).
 - [4] V. Rubakov, M. Sazhin and A. Veryaskin, Phys. Lett. **115B**, 189 (1982).

- [5] R. Fabbri and M.D. Pollock, Phys. Lett. **125B**, 445 (1983).
- [6] A. Starobinski, Sov. Astron. Lett. **9**, 302 (1983).
- [7] L. Abbott and M. Wise, Nucl. Phys. **B244**, 541 (1984).
- [8] L. Abbott and D. Harari, Nucl. Phys. **B264**, 487 (1986).

- [9] M. Gasperini and M. Giovannini, Phys. Rev. **D47**, 1519 (1993).
- [10] R. Brustein, G. Gasperini, G. Giovannini and G. Veneziano, Phys. Lett. **B361**, 45 (1995).
- [11] A. Buonanno, M. Maggiore and C. Ungarelli, Phys. Rev. **D55**, 3330 (1997).
- [12] L.A. Boyle, P.J. Steinhardt and N. Turok, Phys. Rev. **D69**, 127302 (2004).
- [13] B. Boisseau, G. Esposito-Farese, D. Polarski, A.A. Starobinsky, Phys. Rev. Lett. 85, 2236 (2000).
- [14] Y.S. Piao and E Zhou, Phys. Rev. **D68**, 083515 (2003).
- [15] Y.S. Piao and Y.Z. Zhang, Phys. Rev. **D70**, 063513 (2004).
- [16] P.F. Gonzalez-Diaz and J.A. Jimenez-Madrid, Phys. Lett. **B596**, 16 (2004).
- [17] A. Anisimov, E. Babichev and A. Vikman, astro-ph/0504560.
- [18] M. Baldi, F. Finelli and S. Matarrese, astro-ph/0505552.
- [19] S. Nojiri and S.D. Odintsov, hep-th/0506212; hep-th/0507182.
- [20] Y.S. Piao and Y.Z. Zhang, Phys. Rev. **D70**, 043516 (2004); Y.S. Piao, Phys. Lett. **B606** (2005) 245.
- [21] J. Khoury, P.J. Steinhardt and N. Turok, astro-ph/0302012.
- [22] M.S. Turner, J.E. Lidsey and M. White, Phys. Rev. **D48**, 4613 (1993).
- [23] M.S. Turner, Phys. Rev. **D55**, 435 (1997).
- [24] K.S. Thorne, in “300 Years of Gravitation”, edited by S. Hawking and W. Israel.
- [25] B. Allen, gr-qc/9604033.
- [26] A. Buonanno, gr-qc/0303085.
- [27] B. Allen and S. Koranda, Phys. Rev. **D50**, 3713 (1994).